arXiv:1302.4453v1 [cond-mat.quant-gas] 18 Feb 2013

Quantum fragile matter: mechanical excitations of a Reggeon ion chain

Philipp Strack[1,*] and Vincenzo Vitelli[2]
[1]Department of Physics, Harvard University, Cambridge MA 02138
[2]Instituut-Lorentz for Theoretical Physics, Leiden University, Leiden NL 2333 CA, The Netherlands

(Dated: February 20, 2013)

This paper proposes to study *quantum fragile* materials with small linear elasticity and a strong response to zero-point fluctuations. As a first model, we consider a non-unitary (but $\mathcal{PT}$-symmetric) massive quantum chain with a Reggeon-type cubic nonlinearity. At the critical point, the chain supports neither the ordinary quantum phonons of a Luttinger liquid, nor the supersonic solitons that arise in classical fragile critical points in the absence of fluctuations. Quantum fluctuations, approximately captured within a one-loop renormalization group, give rise to mechanical excitations with a nonlinear dispersion relation and dissipative spectral behavior. Models of similar complexity should be realizable with trapped ions.

PACS numbers: 37.10.Ty, 42.65 -k, 43.35.+d, 11.10.Hi

I. INTRODUCTION

Soft materials, such as granular media at vanishing external pressure or random polymer networks with loose connectivity, exhibit large deformations in response to applied mechanical perturbations. In both cases, there is a geometrical or topological control parameter that determines the magnitude of the elastic moduli: the average overlap between grains or the mean coordination number of the network. By tuning these parameters one can reach a mechanical state characterized by a vanishing linear response, often termed sonic vacuum since the linear speed of sound vanishes[1–4]. When this happens even the tiniest strains propagate as supersonic solitons (or shocks) rather than ordinary phonons. The much studied (un)jamming transition is also marked by a progressive loss of rigidity that triggers a strongly non-linear response[5–7]. At the transition these materials are not just soft they become fragile.

The study of the impact of thermal fluctuations on these fragile states is still in its infancy. This is partly due to the fact that granular media and foams, that form the most natural experimental arena to study these phenomena, are simply too large for thermal fluctuations to play any significant role. However, the existence of these athermal critical points rests on fine-tuning geometric parameters of their architecture that would be very sensitive to fluctuations if they were present.

In this paper, we investigate a simple model that, at its critical point, displays a new quantum fragile behavior. As in the known soft counterparts, the mechanical response (the quadratic term in their Hamiltonian) vanishes, but in our model the mechanical properties are strongly affected by zero point fluctuations in a way reminiscent of quantum phase transitions. The notion of fragility that we consider in the present study is not necessarily related to the onset of a plastic response[24,25]. Instead, we denote a classical or quantum mechanical system as fragile, if a parameter can be tuned such that the quadratic part of its Hamiltonian becomes subdominant so that its response to mechanical perturbations is intrinsically non-linear. Recent developments with trapped ions[8–12], Rydberg atoms[13,14] and quantum optomechanics[15,16] give hope that synthesizing soft-matter inspired non-linear systems in the quantum regime may soon be within reach.



FIG. 1: (Color online) Sketch of the model Eq. (1) of a chain of quantum point particles in a local, pinning potential connected by springs. The blue, straight spring describes the harmonic term for the nearest-neighbor strain with strength A. The red, curved spring describes the Reggeon-type cubic nonlinearity for the strain with strength ig. See Ref. 9 for a feasibility study of a related Frenkel-Kontorova chain with trapped ions in optical potentials, Ref. 10 for small-polaron formation with trapped ions, and Ref. 11 for a trapped ion experiment of coupled quantized mechanical oscillators.

The main physical result of our analysis is that the basic excitations at the critical point of a Reggeon ion chain (Fig. 1, Eq. (1)), are neither ordinary phonons, that cannot exist because of the vanishing rigidity, nor the strongly non-linear solitons that characterize similar critical points in the absence of fluctuations. Instead, quantum fluctuations give rise to mechanical excitations with non-linear dispersion relations. This signature can be traced to the emergence of a length-scale dependent sound speed (or equivalently elastic modulus) that we study with the renormalization group to one-loop order.

A. Model

Specifically, we consider a non-unitary but $\mathcal{PT}$-symmetric massive quantum chain with the Hamiltonian operator

$$\hat{H} = \sum_{\ell=1}^{N} \frac{1}{2}\hat{p}_\ell^2 + m^2\,\hat{x}_\ell^2 + \frac{A}{2}\left(\hat{x}_\ell - \hat{x}_{\ell+1}\right)^2 + i\,\frac{g}{36}\left(\hat{x}_\ell - \hat{x}_{\ell+1}\right)^3 \ , \tag{1}$$

describing N point-like quantum particles subject to a local potential of strength m^2 (arising for example in a deep commensurate pinning potential), that interact with their nearestneighbors with a harmonic-chain type coupling A and a cubic nonlinearity with (imaginary) coupling constant ig (see Fig. 1 for a sketch). Here $\hat{x}_\ell$ denotes the deviation from the equilibrium position of the ℓ'th particle and $\hat{p}_\ell$ is the conjugate momentum operator fulfilling the discrete commutation relation: $[\hat{p}_\ell, \hat{x}_{\ell'}] = -i\delta_{\ell\ell'}$ ($\hbar = 1$, lattice constant a=1). The Hamiltonian in Eq. (1) is not Hermitian – it is not invariant under combined matrix transposition and complex conjugation. In addition to translational invariance in space $\ell \rightarrow \ell + n$ with n integer and time, Eq. (1) is invariant under a simultaneous parity transformation ($\mathcal{P}$: $\hat{x} \rightarrow -\hat{x}$) and complex conjugation or time-reversal ($\mathcal{T}$).

Three noteworthy features of Eq. (1) are:

(i) Reggeon-type, non-unitary field theories with cubic nonlinearity have a successful history of describing geometric phenomena such as directed percolation[17–19].

(ii) The cubic nonlinearity ig is chosen proportional to the strain similar to the granular chain[1] but with an integer-valued slightly above the Hertz law for spheres where the interaction scales as $(x_\ell - x_{\ell+1})^\alpha$ with $\alpha = 5/2$.[37]

(iii) The initially non-zero mass from the pinning potential explicitly breaks the continuous shift invariance $\hat{x}_\ell \rightarrow \hat{x}_\ell + \delta$ thereby avoiding the Luttinger liquid fixed point of one dimensional quantum systems[33].[38]

The cubic nonlinearity preserves $\mathcal{PT}$-symmetry and also simplifies the mathematical analysis, for example by giving rise to anomalous dimensions already at the one-loop level. Bender and Boettcher[20,21] demonstrated that certain, non-Hermitian, imaginary Hamiltonians, like the one in Eq. (1), can still describe sensible quantum mechanical ground states with real and positive spectra provided they fulfill a combined parity and time-reversal, $\mathcal{PT}$-symmetry. Such $\mathcal{PT}$-symmetric systems have since been realized experimentally in optical waveguides[22] and it seems possible to engineer periodically structured nonlinearities as well[23]. Related chains of oscillators with Hermitian nonlinearities such as the Fermi-Pasta-Ulam problem[26], quantum Frenkel-Kontorova chains[27,28], chains with disorder[29], or chains under applied forces[30] can also be used as model systems to access the strongly non-linear regime considered in the present study.

We can recast the Hamiltonian operator in Eq. (1) as a Lagrangian density,

$$\mathcal{L} = \frac{1}{2}\Big(\partial_\tau \phi_\ell(\tau)\Big)^2 + m^2\Big(\phi_\ell(\tau)\Big)^2 + \frac{A}{2}\Big(\phi_\ell(\tau) - \phi_{\ell+1}(\tau)\Big)^2 + i\frac{g}{36}\Big(\phi_\ell(\tau) - \phi_{\ell+1}(\tau)\Big)^3 , \tag{2}$$

that defines a lattice quantum field theory for a pseudoscalar ϕ living on a one-dimensional lattice with sites numerated by ℓ and τ is the imaginary time coordinate ranging from 0 to ∞ as appropriate for a zero-temperature quantum field theory. Following Ref. 31, we assume ϕ to transform as a pseudoscalar, that is, ϕ changes sign under space reflection $\mathcal{P}$; then the interaction remains $\mathcal{PT}$-invariant since i changes sign under $\mathcal{T}$.



FIG. 2: (Color online) Phases of the Reggeon quantum chain. The critical point marks the beginning of the fragile regime of small pinning potential (m^2) and harmonic spring constant (A) relative to the nonlinearity (ig).

B. Key results

Our main motivation to study the somewhat artificial model Eq. (2) is that it displays some appealing critical features that may occur also in more complicated and realistic systems. At the same time, it is amendable to an approximate one-loop renormalization group (RG) treatment that yields the phase diagram of Fig. 2. In particular, the RG can flow into the highly nonlinear regime close to the critical point, where violent fluctuations of the ϕ's lead to non-Luttinger liquid criticality for the mechanical excitations in the chain. It is well known that the presence of a continuous symmetry in combination with unitarity inevitably leads to Luttinger liquid behavior of conventional gapless quantum liquids in one dimension[32,33]. Both of these constraints are relaxed in the highly non-linear regime of the non-unitary quantum chain. This is the reason why the imaginary cubic nonlinearity acts as a relevant perturbation to the Luttinger liquid behavior and leads to an unusual power-law scaling of correlators. At the critical point, when the chain becomes fragile, the mechanical excitations disperse as $\omega \sim k^z$ with $z = 0.35$ signaling a strongly momentum-dependent sound velocity. The associated spectral function becomes dissipative and displays a quantum critical continuum that may associated with the spontaneous breakdown of the $\mathcal{PT}$-symmetry.

The rest of the paper is structured as follows. In Subsec. II A, we explain our RG set-up and show how the flow equations yield an interacting fixed point. The new physics associated with this fixed point is presented in detail in Subsec. II B. In Sec. III, we draw our conclusions and point toward promising experimental systems.

II. ONE-LOOP RENORMALIZATION GROUP

In order to perform our RG analysis we write the Lagrangian in a Fourier representation using $\phi_\ell(\tau) = \sum_k \phi_k(\tau) e^{ikx_\ell}$, and analogously for frequencies ω. The lattice action in the thermodynamic limit is obtained as:

$$\Gamma_{\Lambda_0}[\phi] = \int_{\omega,k} \phi_{-k}(-\omega)\left(\frac{Z_{\Lambda_0}}{2}\omega^2 + A_{\Lambda_0}\left(1 - \cos[k]\right) + m^2_{\Lambda_0}\right)\phi_k(\omega) - \frac{g_{\Lambda_0}}{6}\int_{\tau,k_2,k_3} \sin\left[k_2 + k_3\right]\phi_{-(k_2+k_3)}(\tau)\phi_{k_2}(\tau)\phi_{k_3}(\tau) , \tag{3}$$

where we have abbreviated $\int_{\omega,k} = \int_{-\infty}^{\infty}\frac{d\omega}{2\pi}\int_{-\pi}^{\pi}\frac{dk}{2\pi}$ and $\int_{\tau,k_2,k_3} = \int_0^{\infty} d\tau \int_{-\pi}^{\pi}\frac{dk_2}{2\pi}\int_{-\pi}^{\pi}\frac{dk_3}{2\pi}$. We have introduced already here the cutoff scale $\Lambda \in \{\Lambda_0, 0\}$, where Λ_0 is the "ultraviolet" momentum scale of the order of the inverse lattice spacing at which the renormalization parameters Z_{Λ_0}, A_{Λ_0}, $m^2_{\Lambda_0}$ and g_{Λ_0} take their initial values. The endpoint of the RG flow is at $\Lambda \to 0$ and quantum fluctuations on all scales have been integrated into the effective action $\Gamma_\Lambda[\phi]$. We exclude large momentum transfers from the analysis and approximate the trigonometric functions in Eq. (3) by their leading polynomials $1 - \cos[k] \to \frac{k^2}{2} + ...$ and $\sin[k_2 + k_3] \to k_2 + k_3 + ...$ Now, the cubic vertex becomes proportional to the total momentum that flows through it.

Note that the propagator in the quadratic term in the first line of Eq. (3) has a massive phonon dispersion, leading to spectral weight at

$$\omega_{\text{massive}} = \sqrt{\frac{Ak^2 + m^2}{Z}}\ , \tag{4}$$

with Z, A, and m^2 attaining finite values. Throughout the massive phase, the position of the peak will receive finite renormalizations but otherwise the theory will still support well-defined quasi-particles. When approaching the critical point, for vanishing mass $m^2 \to 0$, one immediately notices that a loop expansion of Eq. (3) will suffer from logarithmic and power-law infrared singularities. We will see below that the RG flow to low frequencies and momenta in the presence of interactions gives rise to fractional gradients in space and time characterized by two different anomalous dimensions.

A. RG Scheme

To analyze the physics of these singularities, we employ the (formally exact) RG flow equation for the effective action $\Gamma_\Lambda[\psi]$, the generating functional for one-particle irreducible correlation functions in the form derived by Wetterich[34,35]. We mainly use this method due to its simplicity and versatility to tailor it to the problem at hand. In principle, similar results should be obtainable within an ϵ-expansion or a field-theoretic RG. We add a regulator R_Λ (specified below) to the quadratic part of our bare action Eq. (3) that introduces a cutoff dependence into the effective action so that $\Gamma_\Lambda[\phi]$ smoothly interpolates between the bare action, Eq. (3), at the ultraviolet scale $\Gamma_{\Lambda=\Lambda_0}[\phi] = \Gamma_{\Lambda_0}[\phi]$ and the fully renormalized effective action in the limit of vanishing cutoff: $\lim_{\Lambda\to 0}\Gamma_\Lambda[\phi] = \Gamma[\phi]$. The Wetterich equation

$$\partial_\Lambda \Gamma_\Lambda[\phi] = \frac{1}{2}\mathrm{Tr}\Big[\frac{\partial_\Lambda R_\Lambda}{\Gamma^{(2)}_\Lambda[\phi] + R_\Lambda}\Big] \tag{5}$$

has a one-loop structure and in a vertex expansion the β-functions for the n-point correlators are determined by (cutoff derivatives of) one-particle irreducible one-loop diagrams with fully dressed propagators and vertices. The Tr is here just a frequency and momentum integration and $\Gamma^{(2)}_\Lambda[\phi] = \frac{\partial^2\Gamma_\Lambda[\phi]}{\partial\phi^2}$ is the second functional derivative with respect to the fields ϕ.



FIG. 3: Diagrams for the flow equations.

In the course of the RG flow, new terms, not present in the bare action, will be generated. The objective of the present paper is to choose a simple truncation that reveals the qualitatively new physical features arising from the interplay of quantum fluctuations and the imaginary nonlinearity. We therefore focus on the flow of the scale-dependent phonon frequency renormalization factor Z_Λ, the momentum renormalization factor A_Λ, the mass m^2_Λ and the non-linear coupling g_Λ. No local cubic interaction $ig_{\text{loc}}\phi^3_\ell$ is generated at one-loop. This is because the k-integration in the triangle diagram (see Fig. 3) picks up a factor k^3 and vanishes by symmetry.

As an infrared regulator we use a mass-like cutoff $R_\Lambda = A_\Lambda \Lambda^2$ so that $\partial_\Lambda R_\Lambda = 2A_\Lambda\Lambda$ where, as customary, we drop the (higher-order) term $\sim \partial_\Lambda A_\Lambda$ in the scale-derivative of the cutoff. Now, the infrared singularities at low frequencies and momenta of the phonon propagator

$$G_\Lambda(\omega, k) = \left[\Gamma^{(2)}_\Lambda[\psi] + R_\Lambda\right]^{-1} = \frac{1}{Z_\Lambda\omega^2 + A_\Lambda k^2 + m^2_\Lambda + R_\Lambda} \tag{6}$$

are regulated without by-hand changing the "relativistic" scaling of the bare phonon dispersion $\omega \sim k$. We now plug in Eq. (3) supplemented by the mass and regulator terms into Eq. (5). By comparing coefficients of the fields in a vertex expansion, we obtain the flow equations for Z_Λ, A_Λ, m^2_Λ, and g_Λ. The corresponding diagrams are shown in Fig. 3.

Upon using rescaled variables, the flow equations can be brought into a simple form that facilitates comparison to the well known Wilson-Fisher fixed point of the ϕ^4 theory below its upper critical dimension. The anomalous dimensions absorb singular renormalizations to the frequency (Z) and momentum (A) factor of the phonon propagator, respectively:

$$\eta^Z_\Lambda = -\frac{\Lambda}{Z_\Lambda}\partial_\Lambda Z_\Lambda\ , \qquad \eta^A_\Lambda = -\frac{\Lambda}{A_\Lambda}\partial_\Lambda A_\Lambda\ . \tag{7}$$

Upon rescaling the mass and the cubic interaction as

$$\tilde{m}^2_\Lambda = \frac{m^2_\Lambda}{A_\Lambda\Lambda^2}\ , \qquad \tilde{g}_\Lambda = \frac{g_\Lambda}{\Lambda A^{5/4}_\Lambda Z^{1/4}_\Lambda}\ , \tag{8}$$

we obtain their β-functions:

$$\Lambda\partial_\Lambda \tilde{m}_\Lambda^2 = \left(-2+\eta_\Lambda^A\right)\tilde{m}_\Lambda^2 - \frac{1}{8\pi}\tilde{g}_\Lambda^2\frac{1}{1+\tilde{m}_\Lambda^2} \tag{9}$$

$$\Lambda\partial_\Lambda \tilde{g}_\Lambda = \left(-1+\frac{5}{4}\eta_\Lambda^A+\frac{1}{4}\eta_\Lambda^Z\right)\tilde{g}_\Lambda + \frac{1}{8\pi}\tilde{g}_\Lambda^3\frac{1}{\left(1+\tilde{m}_\Lambda^2\right)^2}\ . \tag{10}$$

The anomalous dimensions for the frequency and momentum factors, respectively, are different

$$\eta_\Lambda^Z = -\frac{1}{48\pi}\tilde{g}_\Lambda^2\frac{1}{\left(1+\tilde{m}_\Lambda^2\right)^2} \tag{11}$$

$$\eta_\Lambda^A = \frac{5}{16\pi}\tilde{g}_\Lambda^2\frac{1}{\left(1+\tilde{m}_\Lambda^2\right)^2}\ . \tag{12}$$

In the infrared $\Lambda \to 0$, these flow equations have two fixed points. A trivial, non-interacting fixed point $\tilde{g}_{\Lambda\to 0} = 0$ with zero anomalous dimensions. More interesting is the infrared-stable, interacting fixed point ($\tilde{g}_* \neq 0$) with finite anomalous dimensions:

$$\eta_*^Z = -0.04\ , \qquad \eta_*^A = 0.61\ . \tag{13}$$

We note that, as usual, the precise numerical values of the critical exponents depend on the renormalization scheme adopted, for example on the choice of the cutoff function.

B. New physics associated with interacting fixed point

From the β-functions Eq. (9-12), we see that the interacting fixed point separates a massive phase from a region of the phase diagram (not studied in the present work and denoted by a dashed line in Fig. 2) where presumably the $\mathcal{PT}$-symmetry is spontaneously broken.

The existence of a non-Gaussian fixed point with finite $\tilde{g}^*$ is ensured by two conditions: (i) the relevant dimensional running leading to the -1 in the first bracket of Eq. (10) and (ii) the positivity of the second term $\sim \tilde{g}^3$. In terms of power counting, (i) is actually similar to the conventional ϕ^4 Wilson-Fisher fixed point in 3 dimensions. (ii) is a direct consequence of the imaginary nature of the cubic interaction in the Lagrangian Eq. (2). The same mechanism turns out to stabilize the real-valuedness of the fixed points of the local $i\phi^3$ theory[31]. Another signature of the non-unitary interaction is a negative anomalous dimension as it was also pointed out in reference[31] and previously in the classic analysis of a ϕ^3 theory by Fisher[36]. We also find a negative η^Z for the frequency renormalization factor here in Eq. (11).

A distinctive feature of the non-unitary quantum chain considered here is that it is not relativistically invariant nor it seem to become so at the critical point, at least not within our one-loop approximation. The cubic vertex depends linearly on spatial momenta (see below Eq. (2)), since it couples neighboring sites. By contrast, it does not depend in the same way on frequencies because the action is completely local in time. While such frequency-dependent interaction will also be generated in the RG flow, they appear only at higher-order in a loop expansion. As a consequence, at the one-loop order considered here, the anomalous dimension for the spatial momenta, η_*^A, is significantly larger and has the opposite sign to η_*^Z. At the critical point, the phonon propagator takes the form:

$$G^*_{\Lambda\to 0}(\omega,k) = \frac{1}{\omega^{2-\eta_*^Z}+k^{2-\eta_*^A}}\ , \tag{14}$$

resulting in a fractional value for the dynamical exponent at the interacting fixed point

$$z_* = 1+\eta_*^Z-\eta_*^A = 0.35\ . \tag{15}$$

This strongly modifies the phonon dispersion at the critical point to

$$\omega_{\text{critical}} \sim k^{z_*} = k^{0.35} \tag{16}$$

and leads to a critical continuum in contrast to the well-defined quasi-particle peak of the massive phase in Eq. (4).

III. CONCLUSION

In this paper, we presented a one-dimensional quantum system with a critical behavior that is different from a Luttinger liquid. To achieve this, we computed the one-loop renormalization group flow of a quantum harmonic chain supplemented by a local mass term and an imaginary cubic nonlinearity. The critical point we found may be viewed as a quantum fragile state of matter with unusual collective excitations and critical exponents. While we have no mathematical proof that the $\mathcal{PT}$-symmetric $i\phi^3$ quantum chain has a well defined quantum mechanical ground state, it seems reasonable to extrapolate the work of Bender *et al.*[31] on the local $i\phi^3$ theory to a one dimensional lattice with translationally-invariant nearest-neighbor interactions.

In the future, one could study local quantum quenches of Eq. (1) in the strongly nonlinear regime to study solitons[1,23] or shocks[4]. We also work toward making synthetic quantum fragile matter accessible in future experiments by scaling up systems that have achieved nonlinear couplings at the single quantum level such as trapped ions[8–12], quantum optomechanical systems[15,16] or optical waveguides[22,23].

Acknowledgments

We thank S. Diehl, E. G. Dalla Torre, M. Lukin, A. M. Turner, P. Zoller, and especially T. Giamarchi, M. Punk, and S. Sachdev for very helpful discussions. This research was supported by the DFG under grant Str 1176/1-1, by the NSF under Grant DMR-1103860, by the Army Research Office Award W911NF-12-1-0227, by the Center for Ultracold Atoms (CUA) and by the Multidisciplinary University Research Initiative (MURI).

---

[*] Electronic address: pstrack@physics.harvard.edu

[1] V. F. Nesterenko, J. Appl. Mech. Tech. Phys. **5**, 733 (1984).

[2] V. F. Nesterenko, *Dynamics of Heterogeneous Materials* (Springer-Verlag, New York, 2001).

[3] L. R. Gómez, A. M. Turner, M. van Hecke, and V. Vitelli, *Phys. Rev. Lett.* **108**, 058001 (2012).

[4] L. R. Gomez, Ari. M. Turner, and V. Vitelli, Phys. Rev. E **86**, 041302 (2012).

[5] C. S. O'Hern, L. E. Silbert, A. J. Liu, and S. R. Nagel, *Phys. Rev. E* **68**, 011306 (2003).

[6] M. van Hecke, *J. Phys.: Condens. Matter* **22**, 033101 (2010).

[7] N. Xu, V. Vitelli, A. J. Liu, and S. R. Nagel, *Europhys. Lett.* **90**, 56001 (2010).

[8] C. F. Roos, T. Monz, K. Kim, M. Riebe, H. Haeffner, D. F. V. James, and R. Blatt, Phys. Rev. A **77** 040302 (R) (2008).

[9] T. Pruttivarasin, M. Ramm, I. Talukdar, A. Kreuter, and H. Haeffner, New Journal of Physics **13**, 075012 (2011).

[10] V. M. Stojanovic, T. Shi, C. Bruder, and J. I. Cirac, Phys. Rev. Lett **109**, 2050501 (2012).

[11] K. R. Brown, C. Ospelkaus, Y. Colombe, A. C. Wilson, D. Leibfried, and D. J. Wineland, Nature **471** 196 (2011).

[12] R. Blatt and C. F. Roos, Nature Physics **8**, 277 (2012).

[13] T. Peyronel, O. Firstenberg, Q. Y. Liang, S. Hofferberth, A. V. Gorshkov, T. Pohl, M. D. Lukin, and V. Vuletic, Nature **488**, 57 (2012).

[14] S. Sevincli, N. Henkel, C. Ates, and T. Pohl, Phys. Rev. Lett. **107**, 153001 (2011).

[15] W. Lechner, S. J. M. Habraken, N. Kiesel, M. Aspelmeyer, and P. Zoller, arXiv:1212.4691 (2012).

[16] T. Ramos, V. Sudhir, K. Stannigel, P. Zoller, and T. J. Kippenberg, arXiv:1302.1855 (2013).

[17] R. C. Brower, M. A. Furman, and M. Moshe, Phys. Lett. B **76**, 213 (1978).

[18] H. Saleur, and D. Sornette, J. Phys. I France **6**, 327 (1996).

[19] H. Hinrichsen, Adv. Phys. **49** 815 (2000).

[20] C. M. Bender and S. Boettcher, Phys. Rev. Lett. **80**, 5243 (1998).

[21] J. Zinn-Justin, and U. D. Jentschura, J. Phys. A: Math. Theor **43**, 425301 (2010).

[22] A. Guo, G. J. Salamo, D. Duchesne, R. Morandotti, M. Volatier-Ravat, V. Aimez, G. A. Siviloglou and D. N. Christodoulides, Phys. Rev. Lett. **103**, 093902 (2009).

[23] M.-A. Miri, A. B. Aceves, T. Kottos, V. Kovanis, and D. N. Christodoulides, Phys. Rev. A **86**, 033801 (2012).

[24] M. E. Cates, J. P. Wittmer, J.-P. Bouchaud, and P. Claudin, Phys. Rev. Lett. **81** 1841 (1998).

[25] M. E. Cates, and M. R. Evans (Eds.), *Soft and fragile matter*, IoP (2000).

[26] G. Gallavotti (Ed.), *The Fermi-Pasta-Ulam Problem: A Status Report*, Lect. Notes Phys. 728 (Springer, Berlin Heidelberg 2008).

[27] O. V. Zhirov, G. Casati, and D. L. Shepelyansky, Phys. Rev. E **67**, 056209 (2003).

[28] O. V. Zhirov, A. S. Pikovsky, and D. L. Shepelyansky, Phys. Rev. E **83**, 016202 (2011).

[29] F. J. Dyson, Phys. Rev. **92**, 1331 (1953).

[30] S. N. Coppersmith and D. S. Fisher, Phys. Rev. A, **38**, 6338 (1988).

[31] C. M. Bender, V. Branchina, and E. Messina, Phys. Rev. D **85**, 085001 (2012).

[32] T. Giamarchi, *Quantum Physics in One Dimension*, CUP (2004).

[33] F. D. M. Haldane, Phys. Rev. Lett **47**, 1840 (1981).

[34] J. Berges, N. Tetradis, and C. Wetterich, Physics Reports **363**, 223 (2002).

[35] P. Strack, and P. Jakubczyk, Phys. Rev. B **80**, 085108 (2009).

[36] M. E. Fisher, Phys. Rev. Lett. **40**, 1610 (1978).

[37] Note that unlike the grains, the interaction considered in Eq. (1) has both an attractive as well as repulsive component.

[38] We thank S. Sachdev for pointing this out.